

Finiteness following from underlying theory: a natural strategy

Jifeng Yang

School of Management, Fudan University, Shanghai, 200433, P. R. China

A tentative proposal is demonstrated that there is a natural strategy to get rid of unphysical (UV) infinities in QFTs if one adopts the modern standard point of view that a fundamental theory that is complete and well-defined in all respects underlies the QFTs. This simple strategy works in principle for any interaction model and space-time dimension. It provides a physical rationality behind the UV divergence and the conventional renormalization programs and improves the latter in several important aspects.

11.10.Ef; 11.10.Gh; 11.90.+t

I. INTRODUCTION

Now, it has become a standard point of view that a fundamental theory (well defined for the extremely high energy end) underlies the present QFTs that are in fact low energy (LE) effective theories for the phenomena in LE ranges [1]. But as far as the author knows, we are still lacking a formulation that can yield finite results in a natural way that fully makes use of the standard point of view. A new strategy is proposed in Ref. [2] that indicates the power of the standard point of view if one uses it appropriately.

To focus on the UV problem, we will assume from now on that there were no unphysical IR singularity in the LE models in our discussions or we have already had an IR regular formulation for the LE QFTs. (We will discuss in our future works about the IR structure's contribution to the whole formulation—it should be included to arrive at a totally satisfying formulation, especially for QCD-like theories where the IR singularity is rather severe and affects the theories' predictions [3]).

It is convenient to employ a generating functional formalism [4] or a path integral formalism to assemble the Green functions for the LE sectors of the underlying theory. It is natural to expect that the well-defined path integral for an effective sector take the following form,

$$Z_{\{\sigma\}}(\{J^i\}) = \int D\mu(\phi_{\{\sigma\}}^i) \exp\{iS(\phi_{\{\sigma\}}^i; \{J_i\}; \{\sigma\})\} \quad (1)$$

where $\{\sigma\}$ are the fundamental parameters (some fundamental constants, probably including the gravitation constant) from the underlying theory and $\{J^i\}$ are the external sources for LE sectors. The 'elementary fields' $(\phi_{\{\sigma\}}^i)$ in the LE sectors are here appended by the underlying parameters to indicate that they are in fact effective ones. It is easy to see that for different LE physics, the LE limit operation may act upon sets of underlying parameters that differ in part.

Within the path integral formalism, we can easily see that in the LE limit (denoted by $\mathbf{L}_{\{\sigma\}}$)

$$Z^0(\{J^i\}; \overline{\cdots}) \equiv \mathbf{L}_{\{\sigma\}} \int D\mu(\phi_{\{\sigma\}}^i) \exp\{iS(\phi_{\{\sigma\}}^i; \{J_i\}; \{\sigma\})\} \quad (2)$$

$$\neq \int D\mu(\phi^i) \exp\{iS(\phi^i; \{J_i\})\}, \quad (3)$$

where the symbols not appended by the underlying parameters refer to the constants and field parameters given by the present QFTs (i.e., the LE limit has been applied on these 'effective' fields or objects). Thus, generally speaking, we can not let the operation of the LE limit cross the summation over the paths or the intermediate states. In other words, the LE limit operation and the summation over intermediate states *do not commute*,

$$\Delta \equiv [\mathbf{L}_{\{\sigma\}}, \sum_{\{paths\}}] \neq 0. \quad (4)$$

If one ignores such a subtle fact, the resulting formulations would be ill defined and needs regularizations that introduce unjustified artificial substitutes of the true underlying structures, which will in turn lead to divergences may appear and subtraction is needed. This is our opinion about the appearance of the unphysical infinities in some conventional QFT formulations.

In the path integral formalism it is immediate to see that the spectra given by the conventional Hamiltonian models would differ from true ones given by the underlying theory, especially in the UV regions. The deviation is signaled by the ill-definedness or UV divergence in the conventional QFTs. More severe UV divergence implies more severe deviation.

What we are trying to present in the following is that if one starts merely with the existence of the underlying theory and $\{\sigma\}$ (without knowing the details), there is a simple strategy to calculate the amplitudes wanted without introducing any *ad hoc* regularization or cutoff that leads to UV divergence. But, due to the lacking of the true underlying structures, there *must remain* in our approach certain ambiguities signaling the missing of the underlying structures, which are to be fixed in principle by phenomenology and experiments.

This paper is organized in the following way: We will first exemplify our strategy in Feynman graph language for the one-loop case in section II. The treatment of the multi-loop cases is given in section III. Then we discuss some nonperturbative examples in section IV to further illustrate our proposal. Finally we summarize our presentation in Section V. We should note in advance that here we do not claim a satisfactory, systematic and final presentation of a new approach, instead we just put forward an alternative strategy for dealing with ill-definedness.

II. HOW CAN UV FINITE RESULTS BE DERIVED

From our discussion in section I, we see that the Hamiltonians (and hence the propagators and vertices) in their present forms are the LE limits of the ones characterized by $\{\sigma\}$. To interest most readers we will exemplify our strategy in the perturbative Feynman graph language (we remind that our strategy is definitely not confined to the perturbative case and we will exhibit the use of the strategy in some nonperturbative cases in section IV). Thus, according to our postulate, the conventional ill-defined (or divergent) Feynman amplitudes (FAs) are results of illegitimate operation order with the LE limit operation and the internal momenta integration. In formula, if the integrand $f(\{Q_i\}, \{p_j\}, \{m_k\})$ (of an ill-defined FA) corresponds to the integrand $\bar{f}(\{Q_i\}, \{p_j\}, \{m_k\}; \{\sigma_l\})$ given by the underlying theory with $\{Q_i\}, \{p_j\}, \{m_k\}$ and $\{\sigma_l\}$ being respectively loop momenta, external momenta, masses and the fundamental parameters in the underlying theory, then

$$\Gamma^0(\{p_j\}, \{m_k\}; \{\bar{c}\}) = \mathbf{L}_{\{\sigma\}} \bar{\Gamma}(\{p_j\}, \{m_k\}; \{\sigma_l\}) = \mathbf{L}_{\{\sigma\}} \int \prod_i d^n Q_i \bar{f}(\{Q_i\}, \{p_j\}, \{m_k\}; \{\sigma_l\}) \quad (5)$$

$$\neq \int \prod_i d^n Q_i \mathbf{L}_{\{\sigma\}} \bar{f}(\{Q_i\}, \{p_j\}, \{m_k\}; \{\sigma_l\}) = \int \prod_i d^n Q_i f(\{Q_i\}, \{p_j\}, \{m_k\}), \quad (6)$$

where $\{\bar{c}\}$ denotes the definite constants arising from the LE limit operation. Γ^0 and $\bar{\Gamma}$ are well-defined (finite) but $\int \prod_i d^n Q_i f(\{Q_i\}, \{p_j\}, \{m_k\})$ is ill defined. That means, the commutator

$$\delta_{\{\sigma\}} = \left[\mathbf{L}_{\{\sigma\}}, \int \prod_i d^n Q_i \right] \quad (7)$$

only vanishes identically for convergent (i.e., well-defined) FAs, otherwise we encounter divergence or ill-definedness in FAs. The deviation of the effective formalism is not detected by the convergent FAs, or these amplitudes are well defined in the present QFTs. This is an extremely important fact for our purpose in the following.

As the underlying theory or the amplitudes $\bar{f}(\dots; \{\sigma_l\})$ are unavailable by now, we have to find a way to approach the truth—the $\Gamma^0(\{p_j\}, \{m_k\})$'s—from our present partial knowledge about the LE sectors of the underlying quantum theory—the present form of the Hamiltonians or Lagrangians as LE sectors of the underlying theory.

Here is our strategy for extracting finite results out of effective formulations of QFT: (1) First we try to perform certain *legitimate* operations (say, Ω) on the objects *so that* the LE limit operation commutes with the summation on the resulted objects if they do not commute on the original objects. (2) Then we can safely perform the intermediate states summation (e.g., loop integration) on the resulted objects. (3) At last, we perform the inverse operations (say, Ω^{-1} , which should also be legitimate) to go back and the final expressions should be (UV) finite by construction but probably ambiguous at the meantime due to our lack of knowledge about the underlying structures. The ambiguities should be fixed from the phenomenological and experimental inputs as mentioned above. The operations (Ω 's) and their inverse are of the main concern in our strategy and they would often be operations with respect to the parameters external to the intermediate states since the objects of interests are expressed in terms of these parameters after all.

The more general framework effecting the natural strategy proposed here will be the subject of our future investigations. Here, we only show how the strategy works in a simple way for the Feynman graph approach, i.e., we exemplify

our simple strategy in the Feynman graph language. First we show that the following important relation holds for 1-loop case ill-defined FAs (c.f. Eq.(5) for 1-loop case)

$$\int d^n Q (\partial_{p_j})^\omega f(Q, \{p_j\}, \{m_k\}) = (\partial_{p_j})^\omega \Gamma^0(\{p_j\}, \{m_k\}), \quad (8)$$

with $\omega - 1$ being the usual superficial divergence degree of $\int d^n Q f(Q, \{p_j\}, \{m_k\})$ so that the left hand side of Eq.(8) exists (finite), $(\partial_{p_j})^\omega$ denoting the differentiation's with respect to the external parameters $\{p_j\}$'s of the amplitude and $\Gamma^0(\dots)$ is the LE limit of the amplitude calculated in the underlying theory (i.e., the internal momentum integration is performed first). It is easy to see that the operation $(\partial_{p_j})^\omega$ leads to convergent graphs or objects.

The proof is very simple, since

$$\begin{aligned} \int d^n Q (\partial_{p_j})^\omega f(Q, \{p_j\}, \{m_k\}) &= \int d^n Q (\partial_{p_j})^\omega \mathbf{L}_{\{\sigma\}} \bar{f}(Q, \{p_j\}, \{m_k\}; \{\sigma_l\}) \\ &= \int d^n Q \mathbf{L}_{\{\sigma\}} (\partial_{p_j})^\omega \bar{f}(Q, \{p_j\}, \{m_k\}; \{\sigma_l\}) = \mathbf{L}_{\{\sigma\}} \int d^n Q (\partial_{p_j})^\omega \bar{f}(Q, \{p_j\}, \{m_k\}; \{\sigma_l\}) \\ &= \mathbf{L}_{\{\sigma\}} (\partial_{p_j})^\omega \bar{\Gamma}(\{p_j\}, \{m_k\}; \{\sigma_l\}) = (\partial_{p_j})^\omega \Gamma^0(\{p_j\}, \{m_k\}). \end{aligned} \quad (9)$$

The second and the fifth steps follow from the commutativity of the two operations $(\partial_{p_j})^\omega$ and $\mathbf{L}_{\{\sigma\}}$ as they act on different arguments, the third step is due to the existence of $\int d^n Q (\partial_{p_j})^\omega f(Q, \dots)$ and the fourth is justified from the existence of $\int d^n Q \bar{f}(Q, \dots; \{\sigma_l\}) (= \bar{\Gamma}(\dots; \{\sigma_l\}))$ by postulate.

It is clear that here the differentiation with respect to the external momenta and its 'inverse'-indefinite integration with respect to the same momenta play the role of the certain operations stated above.

The right hand side of Eq.(8) can be found now as the left end exists as a nonpolynomial (nonlocal) function of external momenta and masses, i.e., denoting it as $\Gamma^0_{(\omega)}$,

$$(\partial_{p_j})^\omega \Gamma^0(\{p_j\}, \{m_k\}) = \Gamma^0_{(\omega)}(\{p_j\}, \{m_k\}). \quad (10)$$

To find $\Gamma^0(\{p_j\}, \{m_k\})$, we integrate both sides of Eq.(10) with respect to the external momenta " ω " times indefinitely to arrive at the following expressions

$$\left(\int_p \right)^\omega [(\partial_p)^\omega \Gamma^0(\{p_j\}, \{m_k\})] = \Gamma^0(\{p_j\}, \{m_k\}) + N^\omega(\{p_j\}, \{c_\omega\}) = \Gamma_{npl}(\{p_j\}, \{m_k\}) + N^\omega(\{p_j\}, \{C_\omega\}) \quad (11)$$

with $\{c_\omega\}$ and $\{C_\omega\}$ being arbitrary constant coefficients of an $\omega - 1$ order polynomial in external momenta N^ω and $\Gamma_{npl}(\{p_j\}, \{m_k\})$ being a definite nonpolynomial function of momenta and masses. Evidently $\Gamma^0(\{p_j\}, \{m_k\})$ is not uniquely determined within conventional QFTs at this stage. That the true expression

$$\Gamma^0(\{p_j\}, \{m_k\}) = \Gamma_{npl}(\{p_j\}, \{m_k\}) + N^\omega(\{p_j\}, \{\bar{c}_\omega\}), \quad \bar{c}_\omega = C_\omega - c_\omega \quad (12)$$

contains a definite polynomial part (unknown yet) implies that it should have come from the LE limit operation on $\bar{\Gamma}(\{p_j\}, \{m_k\}; \{\sigma_l\})$ (see Eq.(5)) as the usual convolution integration can not yield a polynomial part—an indication of the incompleteness of the formalism of the QFTs.

We can take the above procedures as efforts for rectifying the ill-defined FAs and "represent" the FAs with the expressions like the right hand side of Eq.(11), i.e.,

$$\int d^n Q f(Q, \{p_j\}, \{m_k\}) \Rightarrow \Gamma_{npl}(\{p_j\}, \{m_k\}) + N^\omega(\{p_j\}, \{C_\omega\}) \quad (13)$$

with " $\Rightarrow$ " indicating that left hand side is rectified as the right hand side. That the ambiguities reside only in the local part means that the QFTs are quite effective in the LE limit.

To find the $\{\bar{c}_\omega\}$'s in Eq.(12) we need inputs from the physical properties of the system (such as symmetries, invariances and unitarity) and a complete set of data from experiments [5,6] (if we can derive them from the underlying theory all these requirements would be automatically fulfilled). In other words, all the ambiguities should be fixed according to this principle. Similar approach had been adopted by Llewellyn Smith to fix ambiguities on Lagrangian level by imposing high energy symmetry, etc. on relevant quantities [6]. It is also the physical reasoning followed by the conventional renormalization programs.

As we have seen, the $\bar{c}_\omega$'s arise in fact from the low energy limit operation on the objects calculated in the underlying theory, they should be uniquely defined up to possible equivalence. Different or inequivalent choices of these constants simply correspond to different LE theories (amount to being defined by different underlying theories). Since different regularizations and/or renormalization conditions might correspond to inequivalent choices of the constants, (the problem might be more severe in nonperturbative cases [7]), they might lead to different LE theories that even could not faithfully describe relevant low energy physics. Although the underlying structures do not appear explicitly in the LE formulations, they are not totally decoupled from the effective ones and they 'stipulate' the effective sectors indirectly through the constants $\{\bar{c}\}$. In other words, these constants missing from the effective Hamiltonian models are just what we need to define the quantum theory completely in addition to the canonical parameters (masses and couplings). Their missing in the Hamiltonian or Lagrangian level does not mean they do not exist. As we will see in section IV, the former studies on the self-adjoint extension [8] of some quantum mechanical Hamiltonians just support our point of view here.

III. MULTI-LOOP CASE

Since the UV divergence would appear if one first take the limit before doing loop momenta integrations, our strategy here is just to move the limit operator $\mathbf{L}_{\{\sigma\}}$ across the loop integration operations in such a way that no potential divergence is left over.

For any multi-loop graph Γ (we will use the same symbol to denote the graph and the associated FA if it is not confusing), we should again start with the amplitude derived from the underlying theory, i.e., $\bar{\Gamma}(\dots; \{\sigma\})$ with the same graph structure. All the internal lines and vertices are again understood to be given by the underlying theory, characterized by the presence of the parameters $\{\sigma\}$. Then the LE limit of $\bar{\Gamma}(\dots; \{\sigma\})$ produces the definite constants $\{c^0\}$ that are unknown to us yet

$$\Gamma^0(\dots; \{c^0\}) = \mathbf{L}_{\{\sigma\}} \bar{\Gamma}(\dots; \{\sigma\}) = \mathbf{L}_{\{\sigma\}} \int \prod_l d^n l \bar{f}_\Gamma(\{l\}, \dots; \{\sigma\}) \quad (14)$$

where $\bar{f}_\Gamma(\{l\}, \dots; \{\sigma\})$ denotes the integrand obtained from the underlying theory corresponding to the graph Γ and the dots refer to the LE parameters like external momenta, mass parameters and coupling constants. Other symbols are self-evident. We will use in the following $\omega_\gamma - 1$ to denote the overall divergence index [4] for any graph γ and $\{l\}$ to represent the internal momenta and all the partial differentiation operators and their 'inverse' ($\partial_{\omega_\gamma}^{-1}$) act upon the momenta only 'external' to the very internal integration of the graph under consideration. In certain cases, masses are appropriate or efficient external parameters to work with.

If the graph is totally convergent, then Γ^0 contains no UV ambiguity and the limit operation can cross all the internal integrations to act upon the integrand to yield the product of the propagators and vertices given by the present QFTs. But once there is any potential UV ill-definedness associated with any internal integration, one must proceed in the following way (suppose that a graph Γ contains at least an overall divergence)

$$\begin{aligned} \Gamma^0(\dots; \{c_i^0\}) &= \mathbf{L}_{\{\sigma\}} \int \prod d^n l \bar{f}_\Gamma(\{l\}, \dots; \{\sigma\}) \\ \Rightarrow \partial_{\omega_\Gamma}^{-1} \mathbf{L}_{\{\sigma\}} \int \prod d^n l \partial^{\omega_\Gamma} \bar{f}_\Gamma(\{l\}, \dots; \{\sigma\}) &= \sum_{\{\gamma\}=\partial^{\omega_\Gamma} \Gamma} \partial_{\omega_\Gamma}^{-1} \mathbf{L}_{\{\sigma\}} \int \prod d^n l \bar{f}_\gamma(\{l\}, \dots; \{\sigma\}). \end{aligned} \quad (15)$$

The differentiation with respect to the external parameters will give rise to a sum of graphs $\{\gamma\}$ (without overall divergence) from the original graph Γ . (Note that any overlapping divergence will be killed by the differentiation operation, only non-overlapping divergences remain, i.e., the overlapping divergences are disentangled [5]). If there is no more ill-definedness (in any subgraph), one can move the limit operator across all the internal integrations to act directly upon the integrands $\bar{f}_\gamma(\{l\}, \dots; \{\sigma\})$ just like in the totally convergent graph case. Then one can carry out all the loop integrations without any trouble for each graph γ and then sum them up and finally apply the 'inverse' operator with respect to the parameters (usually momenta) external to the graph Γ (and each γ).

Now suppose there are still some ill-defined subgraphs in each γ . In this case, each graph in the set $\partial^{\omega_\Gamma} \Gamma$ can be expressed as a 'product' of disconnected divergent (at least superficially divergent) subgraphs (each subgraph itself may contain overlapping divergences). The LE limit operator crossed all the other parts and stopped before the divergent subgraphs. In formula, for each graph γ , it is

$$\begin{aligned} \partial_{\omega_\Gamma}^{-1} \mathbf{L}_{\{\sigma\}} \int \prod d^n l \bar{f}_\gamma(\{l\}, \dots; \{\sigma\}) &= \partial_{\omega_\Gamma}^{-1} \left\{ \int \prod d^n \bar{l} g_{\gamma/[\gamma']}(\{\bar{l}\}, \dots) \mathbf{L}_{\{\sigma\}} \prod_{\gamma'_j} \int \prod_{i \in \gamma'_j} d^n l'_i \bar{f}_{\gamma:\gamma'_j}(\{l'\}_{\gamma'_j}, \dots; \{\sigma\}) \right\} \\ &= \partial_{\omega_\Gamma}^{-1} \left\{ \int \prod d^n \bar{l} g_{\gamma/[\gamma']}(\{\bar{l}\}, \dots) \prod_{\gamma'_j} [\mathbf{L}_{\{\sigma\}} \bar{\Gamma}_{\gamma:\gamma'_j}(\dots; \{\sigma\})] \right\}, \end{aligned} \quad (16)$$

$$\left(\bigcup_{\gamma'_j} \{l'\}_{\gamma'_j} \right) \bigcup \{\bar{l}\} = \{l\}, \quad [\gamma'] \bigcup \gamma/[\gamma'] = \gamma, \quad [\gamma'] = \prod_j \gamma'_j, \gamma'_j \cap \gamma'_k = 0, \quad j \neq k, \quad (17)$$

where all the dots in the expressions refer to the parameters 'external' to the loop integrations for the subgraphs (i.e., to γ'_j)—they contain the external parameters for the original graph Γ (also for all the graphs in $\{\gamma\}$) and the internal momenta in the set $\{\bar{l}\}$. $\bar{\Gamma}_{\gamma:\gamma'_j}$ refers to amplitude derived from the underlying theory that corresponds to each subgraph γ'_j contained in γ . *Since some loop momenta are 'external' to certain subgraphs, one can not first carry out these loop integrations before the ill-defined subgraphs are treated.* This is in sheer contrast to the totally convergent graphs where the loop integration order does not matter.

As the ill-defined subgraphs in $[\gamma']$ are disconnected with each other, we now treat each of them separately as a new 'total' graph just like what we have done with the total graph Γ starting from Eq.(14). Then we go through the procedures from Eq.(14) to Eq.(17) till we encounter new disconnected and ill-defined subgraphs that are in turn to be treated as before. Finally, we will go to the smallest subgraphs that are completely convergent. Now we can finally remove the LE limit operator to get the integrands totally expressed with propagators and vertices given by the effective theories and we can begin to perform all the loop integrations with insertions of various pairs of $\partial_{\omega_\gamma}^{-1}$ and ∂^{ω_γ} and there will appear a natural order of integration due to our procedures above: the smallest subgraphs are integrated first, then the "larger" subgraphs with their subgraphs done already, and then the still "larger" subgraphs and so on, till all integrations are done. [It is worthwhile to note that at each level of the subgraphs, the loop integrations are guaranteed to be convergent due to Weinberg's theorem [9]].

The resulting expression will be a definite nonlocal functions plus nonlocal ambiguities (due to subgraph ill-definedness) and local ambiguities if Γ is suffering from overall divergence,

$$\Gamma^0(\dots; \{c^0\}) \Rightarrow \Gamma(\dots; \{C\}) = \Gamma_0^{npl}(\dots) + \Gamma_1^{npl}(\dots; \{C'\}) + N^{\omega_\Gamma}(\dots; \{\bar{C}'\}), \quad (18)$$

$$\{C'\} \bigcup \{\bar{C}'\} = \{C\}. \quad (19)$$

Here again we used N^{ω_Γ} to denote the polynomial containing the ambiguities ($\{\bar{C}'\}$) appearing due to the overall divergence. Others are nonlocal functions. Different from the single loop case, there are nonlocal ambiguities in this multiloop graph suffering from subgraph divergences (as evident from our treatment) in addition to the nonlocal definite part and the local ambiguous part. The result we obtained ($\Gamma(\dots; \{C\})$) is not what we are really after ($\Gamma^0(\dots; \{c^0\})$), but that is the best we can do with the present QFT.

Now some remarks are in order:

A. It is evident that overlapping divergences are just automatically resolved in our proposal, because the differentiation operators just remove the overlapping ill-definedness by 'inserting' internal lines and vertices to reduce the overall divergence. Thus one need not worry about them any more. This is the utility derived from the differentiation with respect to external parameters (momenta, masses or other massive parameters that might appear in the LE propagators) [5].

B. The amplitudes given by the underlying theory should be invariant under any linear transformations of the internal integration variables. In our treatment of the ill-defined graphs, since every loop integration actually performed is convergent, these transformations do not alter the results of the loop integrations. Due to the 'inverse' operations, these linear transformations would at most change the polynomial part. But that does not matter at all. This observation implies that one should not worry about the variable shifting and routing of the external momenta that belong to the transformations just described if he has noted the ambiguous part.

B1. An immediate corollary to this observation is that, the chiral anomaly, which is conventionally interpreted as due to the variable shifting in relevant linearly divergent amplitude, must have been due to other definite properties. Otherwise, if it were totally due to the local ambiguities, one can well remove them away by choosing appropriate definitions of the constants (or appropriate renormalization conditions). Our direct calculation shows that [10], one kind of definite rational terms (independent of masses) originated the chiral anomaly. Since they are nonlocal and unambiguous, one can not attribute them simply as UV effects. The trace anomaly is also shown to be originated by such kind of rational terms [10]. To our best knowledge, this nontrivial structure (independent of the UV ambiguities) has never been noted before in the old renormalization framework.

B2. Another utility derived from the observation is that, one can choose the routes of flows of the external momenta to be as simple as possible to make the treatments of an ill-defined multi-loop amplitude as easy as possible. For the single loop cases, sometimes one may only focus on the parts of the amplitude that are really divergent. This may yield fewer ambiguities.

C. With the above deduction, one can easily see that the results of any regularization and/or renormalization scheme can be readily reproduced by corresponding choices of the constants. That is, our proposal can lead to a universal formulation for all the regularization and renormalization schemes at least in the perturbative framework. If one wish to pursue calculation efficiency in the first place, one can choose a regularization scheme that saves labor and then replace all the divergent expressions with an ambiguous polynomial in external parameters to arrive at the same expressions that can be obtained in the procedures described above. Conventionally one has to check whether the symmetry properties of a scheme would affect the main body of a theory and/or works very hard in order to setup a scheme as consistently as possible. Our proposal spares the labor of checking every corner of a scheme and just make use of its efficiency.

For the ambiguities, as we have discussed in the section II, we may first impose some novel symmetries and invariances on the amplitudes to reduce the ambiguities to certain degree, then one has to resort to the experimental physics data, a strategy effectively employed in the conventional renormalization approaches. The W-T identities have served as constraints in the conventional renormalization schemes and saved a lot of labor of calculation. They are just certain kinds of graphical relations among the Feynman graphs. So, in the Feynman graph language, we can derive constraints upon the ambiguities from the graph structures [2].

Moreover, it is worthwhile to note that our strategy works in any kind of models as long as they are physically effective and consistent. That is, it applies efficiently to the unrenormalizable theories as well as the renormalizable ones in the conventional terminology. It would be interesting to integrate our approach with the BV anti-field formalism of QFT [11] that has been used to deal with the unrenormalizable theories quite recently [12]. In contrast, we do not need any counterterm in our strategy for whatever kind of unrenormalizable models simply because we do not encounter any divergence, the only trouble is to fix the ambiguities..

As we have pointed out, the Feynman Amplitudes or the 1PI functions are generally parametrized by more than one constants (I will refer to them as 'agent constants') in addition to the phenomenological ones. If the changes in the radiative constants could be completely compensated by that in the phenomenological ones (which is only possible for rather special kind of models), then we might implement a redefinition invariance of the phenomenological constants or parameters for the FAs like in the RG case. Recently we have rederived the Callan-Symanzik like equations governing the scale transformation behaviors of QFTs without divergence and bare parameters. The influences of the underlying structures effected through the agent constants are clearly shown and these agent constants complete the harmony of the scale transformation of the effective QFTs in the sense discussed in Ref. [13]. Moreover, the new equations improved the conventional ones in quite some aspects.

IV. NONPERTURBATIVE EXAMPLES

It is obvious that our proposal works in principle for any model, whether it is a QFT or not. The key observation that the UV ill-definedness is caused by illegitimate order of 'operations' is valid in both perturbative and the nonperturbative contexts. That is to say, our simple and natural strategy should also be applicable to nonperturbative case.

Recently, the cutoff and Dimensional regularizations are compared in nonperturbative context [7] in the hot topic of applying the idea of effective field theory method (EFT [14]) to LE nuclear physics following Weinberg's suggestion [15]. The framework is a non-relativistic quantum mechanics with Delta-potentials, which is ill defined in the short-distance. According to our discussions above, such ill-definedness means that the effective LE models must have failed in the higher energy end. Then it is illegitimate to simply work with the propagators and vertices (or Green functions and potentials) given by such models. Great care must also be taken with respect to the regularization effects. Thus the inequivalence between the cutoff scheme and Dimensional regularization exhibited in Ref. [7] well evidenced the correctness of our arguments given above.

Now let us try to treat the problem within our proposal. Generally, the Lippmann-Schwinger equation for T -Matrix in the simple two-body problems reads (we follow the notation conventions of Ref. [7])

$$T(p', p; E) = V(p', p) + \int \frac{d^d k}{(2\pi)^d} V(p', k) \frac{1}{E^+ - k^2/(2\mu)} T(k, p; E), \quad (20)$$

where E^+ is $E + i\epsilon$, with E non-negative, and μ denotes the reduced mass in the two-body problem. In our point of view, this equation is not well-defined and should be written as the LE limit of that derived from the well-defined

fundamental underlying theory which is unavailable to us by now. [The underlying parameters will be always denoted as $\{\sigma\}$. For different problems or different LE ranges, the contents may differ.] So in our language, Eq.(20) should be rewritten as

$$T(p', p; E; \{\sigma\}) = V(p', p; \{\sigma\}) + \int \frac{d^d k}{(2\pi)^2} V(p', k; \{\sigma\}) G(E^+ - k^2/(2\mu); \{\sigma\}) T(k, p; E; \{\sigma\}), \quad (21)$$

$$V(p', p) \equiv \mathbf{L}_{\{\sigma\}} V(p', p; \{\sigma\}), \frac{1}{E^+ - k^2/(2\mu)} \equiv \mathbf{L}_{\{\sigma\}} G(E^+ - k^2/(2\mu); \{\sigma\}). \quad (22)$$

Eq.(21) is now well-defined in the underlying theory. Thus Eq.(20) is correct only when there is no UV infinities (there is no IR problem in the following discussions for the Delta-potential problem) so that the LE limit operator can cross the internal momentum integration (summation over intermediate states). Otherwise we have to find a legitimate way to let the LE limit operator cross the internal momenta integration.

In the case of Delta-potential, $V(p', p) = C$, but the $V(\dots; \{\sigma\})$ is generally a nonlocal potential before the LE limit is taken. To be rigorous, we write formally

$$T(p', p; E; \{c^0\}) = C + \mathbf{L}_{\{\sigma\}} \left\{ \int \frac{d^d k}{(2\pi)^d} V(p', k; \{\sigma\}) G(E^+ - k^2/(2\mu); \{\sigma\}) T(p', p; E; \{\sigma\}) \right\}, \quad (23)$$

here it is not generally legitimate to move the $V(\dots; \{\sigma\})$ out of the integration to be directly subject to the LE limit operator—which is exactly what was done in the conventional calculations (with only the propagator regularized)—and it is definitely illegitimate to apply the LE limit operator to all the other objects before the integration is done. Thus, in principle, even when the LE potential is local (of course $V(\dots; \{\sigma\})$ is nonlocal), it might be dangerous to simply reduce Eq.(21) to an algebraic one. Only when the ill-definedness is mainly caused by $1/(E^+ - k^2/(2\mu))$ (i.e., it differs greatly from $G(\dots; \{\sigma\})$ in the UV region where $V(\dots)$ differs less from $V(\dots; \{\sigma\})$), could we pull out the true potential to subject it directly to the action of the LE limit operator. In other words, to put Eq.(21) (a correct formulation for Eq.(20)) or Eq. (23) into an algebraic one requires quite nontrivial properties of the potential and the propagator. To focus on the main point, we temporarily assume this condition is satisfied, then we have the well-defined form of the algebraic equation for the T -matrix (which is now parametrized by the new constants $\{c^0\}$ from the LE limit in addition to E),

$$\frac{1}{T^{on}(E; \{c^0\})} = \frac{1}{C} - I(E; \{c^0\}), \quad (24)$$

with

$$I(E; \{c^0\}) = \mathbf{L}_{\{\sigma\}} \int \frac{d^d k}{(2\pi)^d} G(E^+ - k^2/(2\mu); \{\sigma\}). \quad (25)$$

Now we can employ the technique described in sections II and III to calculate the integrals, i.e., first differentiate $G(E^+ - \dots; \dots)$ with respect to E^+ (which is the 'external' parameter in the integral) for appropriate times, secondly perform the LE limit legitimately and carry out the integral thus obtained, finally do the 'inverse' operation with respect to E and we find the followings (note that here one differentiation with respect to E reduces the divergence degree by two)

$$I_{d,odd}(E; \{c^0\}) \Rightarrow I_{d,o}(E; \{c'\}) = -i \frac{2\mu\pi^{d/2+1}}{\Gamma(d/2)(2\pi)^d} (2\mu E)^{d/2-1} + N^{[(d-1)/2]}(2\mu E; \{c'\}); \quad (26)$$

$$I_{d,even}(E; \{c^0\}) \Rightarrow I_{d,e}(E; \{c'\}) = \frac{2\mu\pi^{d/2}}{\Gamma(d/2)(2\pi)^d} (2\mu E)^{d/2-1} \ln(2\mu E/c'_0) + N^{[d/2]}(2\mu E; \{c'\}) \quad (27)$$

with $\{c'\}$ being arbitrary constants—the ambiguities. These expressions can again be viewed as universal parametrization and compared with that given in cutoff regularization and dimensional regularization schemes (C.f. Ref [7]) with the latter ones as special cases.

In terms of the ambiguous (but *finite*) integrals given by Eq.(26,27), the T -matrix is now parametrized by $\{c'\}$ in addition to E like

$$\frac{1}{T^{on}(E; \{c'\})} = \frac{1}{C} - I_{d,\dots}(E; \{c'\}). \quad (28)$$

Again we need to fix the constants $\{c'\}$ rather than to renormalize the interaction constant C .

It is easy to see that following the normalization condition of Ref. [7], we can reproduce the result derived by Weinberg [15] in two or three dimensional space-time. However, there seems to be no necessary constraints on the phenomenological constant C as it is physical rather than 'bare' in our proposal. Thus, this LE framework à la Weinberg works equally well for both the attractive interactions and the repulsive ones in our approach, contrary to the conclusions that EFT framework failed in the repulsive cases where the LE models are believed to be trivial [16]. The nontriviality of the Delta-potential dynamics has also been investigated by Jackiw [17].

This problem can also be examined from another angle. Jackiw had already pointed out [17] that the Hamiltonians for such models are not automatically Hermitian and need self-adjoint extension. This has already been dealt with by mathematicians in the operator theory and has also been extensively discussed in a number of approaches (please refer to [18] for a comprehensive list of the references). The key point is, in such cases, the self-adjointness of the Hamiltonian is never an automatic property. That is, in contrast to the normal case, the contact potential problem à la Schrödinger equation is UV ill-defined. The resolution of the problem gives rise to a family of self-adjoint extensions of the original Hamiltonian operator parametrized by an additional constant, which upon different choices leads to different or inequivalent (LE) physics [8]. *This additional 'family' parameter is just the constant that will surely be predicted from the LE limit operation in our proposal*, as has been emphasized in section II and III. As a matter of fact, there is an approach that is quite the same as ours in spirit, the one based on resolvent formalism [19] where an important object is defined through an equation in which it appeared in a form differentiated with respect to the 'external' parameter—the resolvent variable (energy). Thus this important object is only defined up to an additional parameter (the family parameter in operator theory approach) which is to be determined by other input, just like in our proposal.

I would like to mention a recent calculation [20] of Higgs masses in nonperturbative context employing our proposal. The results thus obtained are neat and clear, in contrast to that performed within the old renormalization framework (see the references in [20]). Especially, the physical pictures are different from that using the old renormalization, which is easy to see from the discussions above.

One could expect that great ease can be found in employing our proposal or its equivalents (in any form known or unknown) in his/her studies in the nonperturbative contexts and the outcome would be quite different and significant. Moreover, within our approach, those phenomenologically oriented models which are unrenormalizable in the usual renormalization schemes could become quite tamed. One can test it with the NJL model and chiral perturbation theory [21] and with gravity [22]. We will further apply our strategy to more concrete calculations and develop more rigorous framework of this strategy in the future.

V. DISCUSSIONS AND SUMMARY

For a complete representation of the world, we should expect that the underlying theory is also well defined in the IR sector. (Our discussions in the introduction about the spectra are partial as we deliberately omitted the IR issues to focus on the UV structures.) It is conceivable that the phenomenological models give wrong information of the IR end spectra signaled by the unphysical IR infinities.

The underlying theory, 'postulated' here, if exists, should contain all the nontrivial UV and IR structural information lost in the effective theories. Then an interesting scenario dawns upon us: for each effective model dominating certain energy range (say, theory T_{mid}), there should exist two other effective models (or sectors) that are most adjacent to this model from the IR end and UV end respectively (say, T_{IR} and T_{UV}). Then it is imaginable that the phenomenological parameters in T_{IR} and/or T_{UV} would at least quite nontrivially improve the IR and/or UV behaviors of the theory T_{mid} . While on the other hand, the T_{mid} contains what T_{IR} (or T_{UV}) needs to improve its UV (or IR) behaviors. Put it another way, the active and 'elementary' modes or fields in T_{IR} will break up in T_{mid} and give way to the new 'elementary' modes active in T_{mid} . Similarly, the 'elementary' modes in T_{mid} will go 'hibernating' as the energy goes down while 'new' elementary modes 'emerge' to dominate spectra in T_{IR} . The relation between the elementary modes in T_{mid} and T_{UV} is in principle just like that between those in T_{IR} and T_{mid} . Of course, there may be modes active in several successive effective models, some may even be active and stable through all energy levels—the 'fossil' modes or fields we mentioned in the introduction. Evidently, the information about those 'elementary' modes in T_{IR} and T_{UV} missing from T_{mid} (i.e., missing from the effective spectrum given by T_{mid}) can contribute to improve the IR and UV behavior of the latter.

Of course the author does not have a solid idea for the answer right now. Thus we only indicate some interesting but plausible picture about the resolution of the unphysical IR problem or the reformulation of the IR end of a theory suffered from IR problems like QCD.

From our proposal, the 'elementary' commutator for a field and its conjugate, if calculated (or formulated) from the underlying theory, must have been at least a nonlocal function(al) parametrized by the underlying parameters

of the underlying theory and must have been closely related with the gravitational interaction and perhaps new fundamental ones, rather than a highly abstract Dirac delta function containing least information. In a sense, the incompleteness of the present QFTs or their ill-definedness is inherent in the present quantization procedure whose most elementary technical building block is the Dirac delta function (called as distribution by mathematicians) that is *extremely singular and oversimplified in the UV ends*. That the distribution theory works necessarily with test function space or appropriate measure, if viewed from physical angle, is equivalent to that we need more 'fundamental structures' in order for some singular functions to make sense, i.e., a necessity of introducing underlying theory or its artificial substitute-regularization.

In a short summary, we discussed the strategy recently proposed by the author and the important consequences following from it. The proposal could overcome many typical difficulties and shortcomings associated with old regularization and renormalization framework and can be applied in principle to any UV ill-defined quantum theories.

ACKNOWLEDGEMENT

The author is very grateful to Professor F. V. Tkachov for his encouragement and enlightening remarks.

-
- [1] S. Weinberg, *The Quantum Theory of Fields*, Vol I, Ch. XII, Section 3, Cambridge University Press, Cambridge, England, 1995.
 - [2] Jifeng Yang, Report No. hep-th/9708104, hep-th/9807037; 'Can QFT be UV finite as well as effective?', pp202-206 in: Proceedings of the XIth International Conference "Problems of Quantum Field Theory" (Dubna, Russia, July 13-17, 1998), Eds.: B. M. Barbashov, G. V. Efimov and A. V. Efremov. Publishing Department of JINR, Dubna, 1999.
 - [3] I.I. Bigi, *et al*, Phys. Rev. **D 50**, 2234 (1994).
 - [4] N.N. Bogoliubov and D. V. Shirkov, *Introduction to the Theory of Quantized Fields*, 4th edition (Wiley, NY 1980).
 - [5] W. E. Caswell and A. D. Kennedy, Phys. Rev. **D 25**, 392 (1982).
 - [6] C. H. Llewellyn Smith, Phys. Lett. **B 46**, 233 (1973).
 - [7] D. R. Phillips, S. R. Beane and T. D. Cohen, Report No. hep-th/9706070 and references therein.
 - [8] P. Gerbert, Phys. Rev. **D40**, 1346 (1989); M. Reed and B. Simon, *Methods of Modern Mathematical Physics, Vol II*, Ch 10, p145 (Academic Press, New York, 1975).
 - [9] S. Weinberg, Phys. Rev. **118**, 838 (1960).
 - [10] Jifeng Yang and G-j Ni, Acta. Phys. Sinica, **4**, 88 (1995); Jifeng Yang, Report No. hep-th/9801004; G-j Ni and Jifeng Yang, Phys. Lett. **B 393**, 79 (1997).
 - [11] For a review, see, J. Gomis, *et al*, Phys. Rep. **259**, 1 (1995).
 - [12] J. Gomis and S. Weinberg, Nucl. Phys. B **469**, 473 (1996) and references therein.
 - [13] Jifeng Yang, Report No. hep-th/9908111.
 - [14] For reviews, see e.g., D. B. Kaplan, Report No. nucl-th/9506035; or A. V. Manohar, Report No. hep-ph/9606222.
 - [15] S. Weinberg, Phys. Lett. **B 251**, 288 (1990); Nucl. Phys. **B 363**, 1 (1991).
 - [16] M.A.B. Bég and R.C. Furlong, Phys. Rev. **D21**, 1370 (1985); T.D. Cohen, Phys. Rev. **C 55**, 67(1997); D.R. Phillips and T.D. Cohen, Phys. Lett. **B 390**, 7 (1997).
 - [17] R. Jackiw, in *M. A. B. Bég Memorial Volume*, A. Ali and P. Hoodbhoy, eds. (World Scientific, Singapore, 1991), pp25-42.
 - [18] A. Alberverio, *et al*, *Solvable Models in Quantum Mechanics*, (Springer-Verlag, Berlin, 1988) and the extensive references therein.
 - [19] A. Grossmann, R. Hoegh-Krohn and M. Mebkhout, J. Math. Phys. **21**, 2376 (1980).
 - [20] G.-j. Ni, *et al*, Report No. hep-ph/9801264.
 - [21] J. Gasser and H. Leutwyler, Ann. Phys. (NY) **158**, 142 (1984); Nucl. Phys. **150**, 465 (1985).
 - [22] J.F. Donoghue, Phys. Rev. **D 50**, 3874 (1994).